

WEYL EQUATION IN GÖDEL TYPE UNIVERSES

Luis O. Pimentel¹, Abel Camacho and Alfredo Macias²

*Departamento de Física Universidad Autónoma Metropolitana-Iztapalapa,
Apartado Postal 55-534, México D. F., MEXICO.*

Abstract:

The Weyl equation (massless Dirac equation) is studied in a family of metrics of the Gödel type. The field equation is solved exactly for one member of the family.

PACS numbers 04.20q, 04.20.Jb, 04.40+c, 98.80.k

1 Introduction

Field equations for different spins have been studied in the Gödel universe¹⁻⁴ and its extensions^{1,5-7}. Here we consider the massless spinor field equation in a family of homogeneous space-times of the Gödel type^{8,9}. The metric of this class of space-times is the cylindrically symmetric stationary type. We write the metric in the following form,

¹email: lopr@xanum.uam.mx

²email: amac@xanum.uam.mx

$$ds^2 = -(dt + md\phi)^2 + dr^2 + dz^2 + (l - m^2)d\phi^2, \quad (1)$$

where m and l are functions of r , they also satisfy one of the following conditions:

$$D := (l + m^2)^{1/2} = A_1 \exp[ar] + A_2 \exp[-ar], \quad \frac{1}{D} \frac{dm}{dr} = C \quad (2)$$

or

$$D = Ar, \quad \frac{1}{D} \frac{dm}{dr} = C, \quad (3)$$

here A_i and C are arbitrary constants. The whole matter content in the universe described by metric (1) rotates uniformly and intrinsically with local angular velocity:

$$\Omega^\mu = \left(0, 0, 0, \frac{-c}{\sqrt{-g}} \frac{dm}{dr}\right) \quad (4)$$

2 FIELD EQUATION

In order to write the field equation for a massless spin 1/2 field we introduce the tetrad $e^\alpha_\mu(x)$, that satisfies the usual relation

$$g_{\mu\nu}(x) = e^\alpha_\mu(x) e^\beta_\nu(x) \eta_{\alpha\beta}. \quad (5)$$

For the present case we can choose

$$e^0_0(x) = 1, \quad e^1_1(x) = 1, \quad e^0_2(x) = \frac{-m}{D}, \quad e^2_2(x) = \frac{1}{D}, \quad e^3_3(x) = 1 \quad (6)$$

where the $x^0 = t, x^1 = r, x^2 = \phi, x^3 = z$.

The curved Dirac matrices that satisfy

$$\gamma_\mu(x)\gamma_\nu(x) + \gamma_\nu(x)\gamma_\mu(x) = -2g_{\mu\nu}(x), \quad (7)$$

are given by

$$\gamma^0(x) = \tilde{\gamma}^0 - \frac{m}{D}\tilde{\gamma}^2, \quad \gamma^1(x) = \tilde{\gamma}^1, \quad \gamma^2(x) = \frac{1}{D}\tilde{\gamma}_2, \quad \gamma^3(x) = \tilde{\gamma}^3, \quad (8)$$

where the $\tilde{\gamma}^\alpha$ are Dirac matrices in flat Minkowski spacetime.

Weyl equation in curved spacetime is

$$\gamma^\mu \nabla_\mu \psi(x) = 0, \quad (9)$$

$$(1 + \gamma^5)\psi = 0 \quad (10)$$

with

$$\nabla_\mu = \partial_\mu + \Gamma_\mu, \quad \Gamma_\mu = \frac{1}{8}[\tilde{\gamma}^\alpha \tilde{\gamma}^\beta] e_\alpha^\nu e_{\beta\nu;\mu}, \quad (11)$$

and

$$\gamma^5 = \gamma^0 \gamma^1 \gamma^2 \gamma^3. \quad (12)$$

In the case of the metric (1) the spinorial connections are,

$$\begin{aligned} \Gamma_0 &= \frac{m'}{4D}\gamma^2\gamma^1, \quad \Gamma_1 = \frac{m'}{4D}\gamma^0\gamma^2, \\ \Gamma_2 &= \frac{m'}{4}\gamma^0\gamma^1 + \frac{mm' + l'}{4D}\gamma^1\gamma^3, \quad \Gamma_3 = 0 \end{aligned} \quad (13)$$

where the prime means derivative with respect to r . Using the standard representation of the gamma matrices, Dirac equation reduces to

$$\gamma^0\psi_{,0} + \gamma^1\psi_{,1} + \frac{1}{D}\gamma^2\psi_{,2} + \gamma^3\psi_{,3} - \frac{m}{D}\gamma^2\psi_{,0} - \frac{m'}{4D}\gamma^2\gamma^0\gamma^1\psi + \frac{2D'}{D}\gamma^1\psi = 0, \quad (14)$$

in the above equation we can be separate variables according to

$$\psi_{\mathbf{k}} = \exp[-i(k_0t + k_2\phi + k_3z)] \begin{pmatrix} \xi_1(r) \\ \xi_2(r) \end{pmatrix} \quad (15)$$

with ξ_i two-component spinors, and k_i constants. The chirality condition, Eq.(10) implies that

$$\xi_1 = \xi_2 = \begin{pmatrix} R_1(r) \\ R_2(r) \end{pmatrix} \quad (16)$$

and Eq.(9) is now

$$R_{2,1} + \frac{1}{D}(k_0m - k_2)R_2 + i\left(\frac{m'}{4D} - k_0 - k_3\right)R_1 = 0, \quad (17)$$

$$R_{1,1} - \frac{1}{D}(k_0m - k_2)R_1 + i\left(k_3 - k_0 - \frac{m'}{4D}\right)R_2 = 0. \quad (18)$$

In all the previous equations we have not used an explicit form for the functions $l(r)$ and $m(r)$, we have been able to solve Dirac equation for the second form of those functions (Eq.(3)). If we eliminate R_2 between the above two equations and using the explicit expression for $m = m_0 + \frac{CA}{2}r^2$ we obtain

$$R_1'' + \left\{ \frac{1}{Ar^2}(\tilde{k}_0r^2 - \tilde{k}_2)[1 - \frac{1}{A}(\tilde{k}_0r^2 - \tilde{k}_2)] + E \right\} R_1 = 0, \quad (19)$$

where we defined

$$\tilde{k}_0 = \frac{k_0AC}{2}, \tilde{k}_2 = k_0m_0 - k_2, E = (k_3 - k_0 - \frac{C}{4})\left(\frac{C}{4} - k_0 - k_3\right) - \frac{2\tilde{k}_0}{A} \quad (20)$$

3 EXACT SOLUTIONS

The solution to Eq.(19) is

$$R_1(r) = (\sqrt{2}\tilde{A}r^2)^{\frac{-1}{4}}[A_1M_{\kappa,\mu}(\sqrt{2}\tilde{A}r^2) + A_2W_{\kappa,\mu}(\sqrt{2}\tilde{A}r^2)], \quad (21)$$

where

$$\tilde{A} = \left(\frac{k_0}{A}\right), \quad (22)$$

and $M_{\kappa,\mu}$ and $W_{\kappa,\mu}$ are the Whittaker functions defined in terms of the confluent hypergeometric functions M and U,

$$M_{\kappa,\mu}(x) = e^{-x/2}x^{1/2+\mu}M\left(\frac{1}{2} + \mu - \kappa, 1 + 2\mu, x\right), \quad (23)$$

$$W_{\kappa,\mu}(x) = e^{-x/2}x^{1/2+\mu}U\left(\frac{1}{2} + \mu - \kappa, 1 + 2\mu, x\right), \quad (24)$$

and the parameters κ and μ in this case are given by

$$\kappa = \frac{E}{\sqrt{8}\tilde{A}}, \quad \mu = \sqrt{\frac{1}{16} + \frac{1}{2}\frac{1}{A}\tilde{k}_2\left(1 + \frac{1}{A}k_2\right)} \quad (25)$$

On this solution we impose the condition that the Whittaker functions must be bounded for all values of r . This condition is realized if and only if

$$\frac{1}{2} + \mu - \kappa = -n \quad (26)$$

noi here n is a positive integer or zer. From the above definitions that is equivalent to

$$\frac{1}{2} + \sqrt{\frac{1}{16} + \frac{1}{2}\frac{1}{A}\tilde{k}_2\left(1 + \frac{1}{A}k_2\right)} - \frac{E}{\sqrt{8}\tilde{A}} = -n, \quad (27)$$

where n is a natural number or zero.

In this way we have been able to solve the Weyl equations for a family of spacetimes of the Gödel type.

4 ACKNOWLEDGMENTS

This work was partially supported by the CONACYT GRANTS 1861-E9212, 3544-E9311.

References

- [1] Cohen J M, Vishveshwara C V and Dhurandhar S V 1980 J. Phys. A, **13** 933.
- [2] Hiscock W A 1978 Phys.Rev., **D17**, 1497.
- [3] Mashoon B 1975 Phys.Rev., **D11**, 2679.
- [4] Pimentel L O and Macias A 1986, Phys. Lett. **A 117**, 325.
- [5] Abd-Eltwab Y A 1993 Il Nuovo Cimento **108 B** 464.
- [6] Soares I D 1981 Phys.Rev., **D23**, 272.
- [7] Villalba V 1993 Modern Physics Letters **8**, 3011.
- [8] Raychaudhuri A K and Thakurta S N G 1980 Phys.Rev., **D22**, 802.
- [9] Chakraborty S K and Bandyopadhyay N 1983 J. Math. Phys., **D24**, 129.